

**The 65th William Lowell Putnam Mathematical Competition**  
**Saturday, December 4, 2004**

A-1 Basketball star Shanille O'Keal's team statistician keeps track of the number,  $S(N)$ , of successful free throws she has made in her first  $N$  attempts of the season. Early in the season,  $S(N)$  was less than 80% of  $N$ , but by the end of the season,  $S(N)$  was more than 80% of  $N$ . Was there necessarily a moment in between when  $S(N)$  was exactly 80% of  $N$ ?

A-2 For  $i = 1, 2$  let  $T_i$  be a triangle with side lengths  $a_i, b_i, c_i$ , and area  $A_i$ . Suppose that  $a_1 \leq a_2, b_1 \leq b_2, c_1 \leq c_2$ , and that  $T_2$  is an acute triangle. Does it follow that  $A_1 \leq A_2$ ?

A-3 Define a sequence  $\{u_n\}_{n=0}^\infty$  by  $u_0 = u_1 = u_2 = 1$ , and thereafter by the condition that

$$\det \begin{pmatrix} u_n & u_{n+1} \\ u_{n+2} & u_{n+3} \end{pmatrix} = n!$$

for all  $n \geq 0$ . Show that  $u_n$  is an integer for all  $n$ . (By convention,  $0! = 1$ .)

A-4 Show that for any positive integer  $n$ , there is an integer  $N$  such that the product  $x_1 x_2 \cdots x_n$  can be expressed identically in the form

$$x_1 x_2 \cdots x_n = \sum_{i=1}^N c_i (a_{i1} x_1 + a_{i2} x_2 + \cdots + a_{in} x_n)^n$$

where the  $c_i$  are rational numbers and each  $a_{ij}$  is one of the numbers  $-1, 0, 1$ .

A-5 An  $m \times n$  checkerboard is colored randomly: each square is independently assigned red or black with probability  $1/2$ . We say that two squares,  $p$  and  $q$ , are in the same connected monochromatic component if there is a sequence of squares, all of the same color, starting at  $p$  and ending at  $q$ , in which successive squares in the sequence share a common side. Show that the expected number of connected monochromatic regions is greater than  $mn/8$ .

A-6 Suppose that  $f(x, y)$  is a continuous real-valued function on the unit square  $0 \leq x \leq 1, 0 \leq y \leq 1$ . Show that

$$\begin{aligned} & \int_0^1 \left( \int_0^1 f(x, y) dx \right)^2 dy + \int_0^1 \left( \int_0^1 f(x, y) dy \right)^2 dx \\ & \leq \left( \int_0^1 \int_0^1 f(x, y) dx dy \right)^2 + \int_0^1 \int_0^1 (f(x, y))^2 dx dy. \end{aligned}$$

B-1 Let  $P(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_0$  be a polynomial with integer coefficients. Suppose that  $r$  is a rational number such that  $P(r) = 0$ . Show that the  $n$  numbers

$$c_n r, c_n r^2 + c_{n-1} r, c_n r^3 + c_{n-1} r^2 + c_{n-2} r, \\ \dots, c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r$$

are integers.

B-2 Let  $m$  and  $n$  be positive integers. Show that

$$\frac{(m+n)!}{(m+n)^{m+n}} < \frac{m! n!}{m^m n^n}.$$

B-3 Determine all real numbers  $a > 0$  for which there exists a nonnegative continuous function  $f(x)$  defined on  $[0, a]$  with the property that the region

$$R = \{(x, y); 0 \leq x \leq a, 0 \leq y \leq f(x)\}$$

has perimeter  $k$  units and area  $k$  square units for some real number  $k$ .

B-4 Let  $n$  be a positive integer,  $n \geq 2$ , and put  $\theta = 2\pi/n$ . Define points  $P_k = (k, 0)$  in the  $xy$ -plane, for  $k = 1, 2, \dots, n$ . Let  $R_k$  be the map that rotates the plane counterclockwise by the angle  $\theta$  about the point  $P_k$ . Let  $R$  denote the map obtained by applying, in order,  $R_1$ , then  $R_2, \dots$ , then  $R_n$ . For an arbitrary point  $(x, y)$ , find, and simplify, the coordinates of  $R(x, y)$ .

B-5 Evaluate

$$\lim_{x \rightarrow 1^-} \prod_{n=0}^{\infty} \left( \frac{1+x^{n+1}}{1+x^n} \right)^{x^n}.$$

B-6 Let  $\mathcal{A}$  be a non-empty set of positive integers, and let  $N(x)$  denote the number of elements of  $\mathcal{A}$  not exceeding  $x$ . Let  $\mathcal{B}$  denote the set of positive integers  $b$  that can be written in the form  $b = a - a'$  with  $a \in \mathcal{A}$  and  $a' \in \mathcal{A}$ . Let  $b_1 < b_2 < \cdots$  be the members of  $\mathcal{B}$ , listed in increasing order. Show that if the sequence  $b_{i+1} - b_i$  is unbounded, then

$$\lim_{x \rightarrow \infty} N(x)/x = 0.$$